


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/68		A1	(11) International Publication Number: WO 99/30163
			(43) International Publication Date: 17 June 1999 (17.06.99)
(21) International Application Number: PCT/IL98/00592 (22) International Filing Date: 7 December 1998 (07.12.98) (30) Priority Data: 122490 7 December 1997 (07.12.97) IL (71) Applicant (for all designated States except US): YEDA RESEARCH AND DEVELOPMENT CO. LTD. [IL/IL]; Weizmann Institute of Science, P.O. Box 95, 76100 Rehovot (IL). (72) Inventors; and (75) Inventors/Applicants (for US only): SHINITZKY, Meir [IL/IL]; Derech Haganim Street 20, 46910 Kfar Shmaryahu (IL). DECKMANN, Michael [DE/FR]; 24, rue de Kreutzberger, F-68500 Guebwiller (FR). (74) Agent: REINHOLD COHN AND PARTNER; P.O. Box 4060, 61040 Tel Aviv (IL).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SKIN TEST FOR SCHIZOPHRENIA			
(57) Abstract <p>A diagnostic method for assaying schizophrenia in a subject is provided wherein a preparation comprising platelet derived proteins or fractions thereof having a pI above about 6.5 is injected into a subject and the occurrence of delayed type hypersensitivity (DTH) reaction at the site of the injection is determined. A positive DTH reaction indicates that the tested subject has a high likelihood of being schizophrenic. The protein preparation used in the diagnostic method is also provided as well as a method for its preparation and a kit for use in the diagnosis of schizophrenia using the above method.</p>			
<div style="display: flex; align-items: center;"><div style="margin-right: 20px;"><p>P₂</p><p>P₁</p><p>A</p><p>C</p></div></div>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SKIN TEST FOR SCHIZOPHRENIA

FIELD OF THE INVENTION

The present invention relates to diagnostic tests, and more specifically to diagnostic tests for psychiatric diseases.

5 LIST OF REFERENCES

The following is a list of references which is considered to be pertinent for the purpose of understanding the background of the invention:

1. Nyland, H., Naess, A. and Lundre, H.: *Lymphocyte subpopulations in peripheral blood from schizophrenic patients. Acta Psychiat. Scand.* 1980; 61; 313-318.
2. Hirata-Hibi, M., Higashi, S., Tachibana, T. and Watanabe, N., *Stimulated lymphocytes in schizophrenia Arch. Gen. Psychiat.* 1982; 39; 82-87.
- 15 3. Coeffey, C.E., Sullivan, J.L. and Rice, R.R., *T lymphocytes in schizophrenia. Biol. Psychiatry* 1983; 18; 113-119.
4. Kolyaskina, G.I. *Blood lymphocytes in schizophrenia - immunological and virological aspects. Adv. Biol. Psychiat.* 1983; 12; 142-149.
5. Bessler, H., Eviatar J., Meshulan. M., Tyano. S., Djaldetti, M. and
20 Sirota P. *Theophyllin-sensitive T-lymphocyte subpopulation in schizophrenic patients. Biol. Psychiatr.* 1987; 22; 1025-1029.
6. Muller, N., Ackenheil, M., Eckstein, R., Hofschuster, E. and Mempel,

- W. *Reduced suppressor cell function in psychiatric patients. Ann. N.Y. Acad. Sci. 1987; **396**; 686-690.*
7. Mihalovic, L.J. and Jankovic B.D. *Effects of intraventricularly injected anti-caudatus antibody on the electrical activity of the cat brain. Nature 1961; **192**; 665.*
8. Rapport, M.M., Karplak, S.E. and Mahadik, S.P. *Biological activities of antibodies injected into the brain. Fed. Proc. 1979; **38**; 2391.*
9. Vartanian, M.E. Doyskina, G.S. Lozovsky, D.V., Burbaera, G.S. and Ignaton, S.A. *Aspects of humoral and cellular immunity in schizophrenia. In: Birth Defects. Original Article Series D. Bergsma and A. Goldstein, eds. vol 14; 339-364; Alan R. Liss; New York, N.Y; 1978.*
10. Rotman, A. *Blood platelets in psychopharmacological research. Prog. Neuropsychopharmacol. 1983; **6**; 135-151.*
11. Pletscher, A. *Biological Psychiatry*, Gea Racagni, ed; Elsevier Science Publisher; 1991; **2**, 354-356.
12. Shinitzky, M., Deckmann, M., Kessler, A., Sirota, P., Rabbs, A. and Elizur, A., *Platelet autoantibodies in dementia and schizophrenia - possible implications for mental disorders. Ann. N.Y. Acad. Sci. 1991; **621**; 205-217.*
13. Kessler, A. and Shinitzky, M., *Platelets from schizophrenic patients bear autoantibodies that inhibit dopamine uptake. Psychobiology 1993; **21**; 229-306.*
14. PCT Patent Application WO 95/23970.
15. Deckmann *et al.*, *Italian Journal of Psychiatry and Behavioural Sciences*, **6**:29-34, 1996.

The above references will be acknowledged in the text below by indicating their number from the above list shown in brackets.

BACKGROUND OF THE INVENTION

Schizophrenia is a syndrome which encompasses a variety of symptoms including paranoia, auditory hallucination, delusions, catatonia, bizarre behavior and emotional withdrawal. Schizophrenia affects about 1% of the total population. Its economical and social burden on society is enormous since onset occurs in youth thus requiring patients to be under medical and psychiatric supervision for most of their lives. Schizophrenia is therefore one of the most costly diseases in the industrialized world.

Since the biochemical basis of schizophrenia has not yet been elucidated, diagnosis today is still based solely upon psychiatric evaluation. Furthermore, no therapy is currently available for schizophrenia although the symptoms may be ameliorated by neuroleptic drugs.

Many reports have implicated the immune system in the etiology and course of several mental disorders. Serum antibodies which cross-react with brain antigens have been found in the blood of schizophrenic patients⁽¹⁻⁶⁾, thus indicating that schizophrenia is also an autoimmune disease⁽⁷⁻⁹⁾. Furthermore, platelets have been used as a model for neuronal tissue^(10,11) and elevated levels of autoantibodies to platelets have been detected in schizophrenic and demented patients, but not in patients suffering from manic-depressive disorder, depression, personality disorders or schizoaffective disorders⁽¹²⁻¹⁴⁾.

A cellular response against autologous platelets was also demonstrated in schizophrenia patients who showed a delayed type hypersensitivity (DTH) reaction when injected with platelets collected from their own blood⁽¹⁵⁾.

It is therefore the object of the present invention to provide a test for the diagnosis of schizophrenia in a subject.

SUMMARY OF THE INVENTION

A novel finding in accordance with the invention is that schizophrenic patients exhibit a reaction which displays characteristics of a typical delayed-type hypersensitivity reaction when injected with a specific, novel fraction of platelet proteins. This novel fraction is referred to herein as "Pool 2 proteins". The Pool 2 proteins, which constitute an aspect of the invention, are characterized by having an isoelectric point (pI) which is above about 6.5 and preferably within the range of about 6.5 to about 9.5. When injected with Pool 2 proteins, schizophrenic patients show such a delayed-type hypersensitivity (DTH) reaction.

It is to be noted that although the reaction of the patients to the Pool 2 proteins is referred to above and below as a DTH reaction, there may be cases in which the time profile of the reaction of the patient to the Pool 2 proteins will be different than the time profile of a typical DTH reaction.

The present invention thus provides a protein preparation (hereinafter "*Pool 2 proteins*") comprising platelet derived proteins or fractions thereof having an isoelectric point (pI) above about 6.5 and preferably within the range of above 6.5 to about 9.5, said preparation capable of eliciting a DTH reaction in a schizophrenic individual upon injection thereof to the individual.

The present invention further provides a diagnostic method for assaying schizophrenia in a subject comprising:

- (a) obtaining a preparation comprising, as an active component, platelet derived proteins or fractions thereof having an isoelectric point (pI) above about 6.5 and preferably within the range of about 6.5 to about 9.5 ("*pool 2 proteins*");
- (b) injecting said protein preparation into a subject; and
- (c) examining the subject for the occurrence of a delayed type hypersensitivity reaction at the site of the injection, a positive result being a

reaction above that which is observed in non-schizophrenic subjects, indicating that the subject has a high likelihood of being schizophrenic.

In accordance with the invention it was also surprisingly found that the Pool 2 proteins may be either prepared from autologous platelets of the individual to be tested or, alternatively, from a pool of heterologous platelets which were obtained from a number of individuals other than the tested subject. It was also surprisingly found that, in most cases, the DTH reaction in a schizophrenic patient is substantially higher when the individual is injected with such Pool 2 proteins obtained from a pool of heterologous platelets as compared to a lower DTH reaction in the same tested individual injected with Pool 2 proteins prepared from his own autologous platelets. This finding provides the advantage of preparing a Pool 2 protein preparation which may then either be used immediately or alternatively, be stored at appropriate conditions (e.g. refrigeration) and used for various periods of time to diagnose schizophrenia in a large number of individuals. This obviates the need to repeatedly obtain a blood sample comprising platelets from the tested individual immediately prior to carrying out the diagnostic assay of the invention.

Thus, by a preferred embodiment, the present invention further provides a method for the preparation of a reagent for use in diagnosis of schizophrenia, comprising:

- (a) obtaining blood samples from a number of individuals, preparing a pool from said samples and collecting platelets therefrom;
- (b) preparing a protein fraction from said platelet preparation comprising proteins or fractions thereof having a pI of above about 6.5, preferably within the range of about 6.5 to about 9.5.

The Pool 2 proteins prepared from heterologous individuals may either be prepared from a number of individuals suffering from schizophrenia or, alternatively, also from a mixture of blood samples obtained from schizophrenic patients as well as healthy individuals.

5 The present invention yet further provides a diagnostic method for assaying schizophrenia in a subject comprising:

(a) obtaining a blood sample from a number of schizophrenic and/or non schizophrenic individuals other than the tested subject and collecting platelets therefrom;

10 (b) preparing a protein fraction from said platelet separation comprising proteins or fractions thereof having a pI of above about 6.5, preferably within the range of above 6.5 to about 9.5;

(c) injecting said protein preparation into a subject; and

(d) examining the subject for the occurrence of a delayed type
15 hypersensitivity reaction at the site of the injection, a positive result being a reaction above that which is observed in non-schizophrenic subjects, indicating that the subject has a high likelihood of being schizophrenic.

By a preferred embodiment, a method for diagnosis of schizophrenia in an individual is provided as above wherein the protein
20 fraction collected in stage (b) is then stored at appropriate conditions for repetitive use as described in stages (c) and (d) above at later periods of time for a number of tested individuals.

Although, as explained above, there is an advantage in preparing a pool of Pool 2 proteins from a number of individuals other than
25 the individual to be tested, in accordance with the diagnostic method of the invention, it is also possible to use a Pool 2 preparation prepared from autologous platelets of the individual to be tested.

The present invention thus provides a diagnostic method for assaying schizophrenia in a subject comprising;

- (a) obtaining a blood sample from an individual and collecting platelets therefrom;
- 5 (b) collecting proteins or fractions thereof from said platelet sample, said proteins or fractions having a pI of above about 6.5, preferably within the range of about 6.5 to about 9.5;
- (c) injecting said collected proteins or fractions thereof to the tested individual; and
- 10 (d) examining the subject for the occurrence of delayed type hypersensitivity reaction at the site of the injection, a positive result being a reaction above that which is observed in non-schizophrenic subjects, indicating that the subject has a high likelihood of being schizophrenic.

Typically, the Pool 2 proteins will be collected by subjecting
15 the collected platelet proteins to isoelectric focusing wherein the proteins having a pI of above about 6.5, preferably within the range of 6.5 to 9.5 are collected by methods known in the art.

However, in accordance with the invention it has also been found that the Pool 2 protein preparation may also be prepared by methods
20 which do not include isoelectric focusing such as, for example, by extracting the proteins from the platelet sample using for example a detergent, such extraction resulting in Pool 2 proteins having the desired pI values and capable of eliciting a DTH reaction in a schizophrenic patient.

If desired, the Pool 2 protein preparation obtained by any one of
25 the methods mentioned above may be subject to further fractionation steps, e.g. by thin layer chromatography, by high pressure liquid chromatography or by many other purification fractionation methods known, *per se*. Fractions thus obtained can each then be tested for activity, namely for its ability to cause the DTH reaction in schizophrenic patients. Such purified fractions, as

well as individual proteins, polypeptides or peptides among the Pool 2 proteins which are active in eliciting the DTH reaction in schizophrenic patients, are also an aspect of the invention.

The Pool 2 protein preparation prepared in accordance with the invention and used in the diagnostic methods of the invention include proteins purified from the Pool 2 proteins, polypeptides or peptides comprising sequences of such proteins, fractions thereof, as well as proteins, polypeptides or peptides obtained by synthesis or by genetic engineering having a sequence identical or homologous (e.g. above 70%, preferably above 80%, and typically above 90% homology) to that of the proteins of the Pool 2 proteins.

In accordance with the invention, Pool 2 proteins used in the diagnostic assay of the invention are such which are capable of eliciting DTH activity in an injected individual, the DTH activity being tested by the test known in the art. In short, the Pool 2 proteins are intradermally injected into the tested individual at the forearm or thigh and the reaction at the injection site is evaluated after 24, 48 and 72 hours by measuring the reaction diameter around the induration. As mentioned above, there may be cases in which the time profile of the reaction will differ from the typical time profile of a DTH reaction.

The present invention further provides a kit for use in diagnosis of schizophrenia, comprising said Pool 2 proteins, active protein fractions obtained therefrom, or individual active proteins or peptides, derived from said Pool 2 proteins. Preferably, such proteins are provided in either injectable form or in a form suitable for preparing an injectable formulation, e.g. a lyophilysate. Typically, the kit will be provided with instructions for use or a chart or pictures for guidance of the manner of scoring the results.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows the reaction on the forearm of a schizophrenic patient which was injected with the following preparations:

- i. Pool 2 proteins (marked as P2)
- 5 ii. Pool 1 proteins (marked as P1)
- iii. Autologous platelets (marked A)
- iv. PBS (marked as C = control).

The invention will now be illustrated in the following examples, which are annexed to the above drawings.

10

EXAMPLES

A study on the reaction of a subject to an injection of platelets collected from his or her own blood or an injection of Pool 2 proteins was carried out in Israel at The Hospital for Mental Health in Sha'ar Menashe, The
15 Geriatric Hospital in Pardes Hana and at the Weizmann Institute of Science.

Example 1 Injection of autologous proteins

1. Preparation of platelets

10 ml venous blood was collected from a subject and
20 centrifuged for 20 mins. at 20°C and 150 g. The supernatant containing platelets was collected and centrifuged three times for 10 min. at 20°C and 2000 g and the platelets resuspended in phosphate buffered saline (PBS) containing 5 mM EDTA. After the last washing, the platelets were resuspended in sterile PBS at a final concentration of 2×10^8 platelets/ml.

25

2. Autologous skin test and measurement of delayed type hypersensitivity (DTH) reaction

0.1 ml of the platelet suspension obtained as above from a
30 particular subject was injected intradermally back into the forearm of the

same subject. A second injection of 0.1 ml PBS spaced about 10 cm from the point of sample injection served as a control.

DTH reaction at the injection sites was measured in accordance with methods known in the art (Skornik, Y., *et al.*, *Cancer Immunol. Immunother.* 11:93-96, 1981).

Results

The tests were carried out for groups subject:

- a) 18 healthy subjects under the age of 65.
- 10 b) 10 healthy subjects above the age of 65.
- c) 41 schizophrenic patients.
- d) 21 demented patients.

The results of this test are shown in the following Table 1.

Table 1**Autologous (Self-Self) Skin Reaction Against Platelets in Humans
for the Detection of Schizophrenia****Summary of a Multi-Center Study**

Schizophrenic Persons n = 41	Non-schizophrenic Persons n = 49
Skin Reaction positive: 25	Skin Reaction positive: 0
Skin Reaction borderline: 13	Skin Reaction borderline: 1
Skin Reaction negative: 3	Skin Reaction negative: 48
Total Sensitivity: 92%	Total Specificity: 98%

As can be seen, 38 out of the 41 schizophrenic patients showed a DTH reaction, while only one healthy individual under the age of 65 out of the 18 which were tested, reacted positively. Positive reactions were not observed in 10 healthy individuals over the age of 65, as well as in 21 demented patients. Sensitivity of this test is thus 92% and the specificity is 98%.

As seen in the Table, while almost all of the tested schizophrenic patients showed a DTH reaction to the Pool 2 proteins prepared from their autologous platelets, only one healthy individual had a positive DTH reaction. Thus, the diagnostic assay of the invention showed a very high sensitivity and specificity for the diagnosis of schizophrenia in a tested individual.

Example 2 Preparation of Pool 1 and Pool 2 proteins**Methods**

Platelet suspension containing about 20 gr total protein was obtained as in Example 1, and the platelets solubilized with 40 ml of a solution containing 0.5% of the detergents NP-40 and Triton-X-100 for 5 mins. at room temperature with gentle shaking. The solution was then centrifuged at 4000 g for 15 mins. at 20°C. The supernatant was collected, and the pellet was subjected to two further extractions with 10 ml 0.1% Triton-X-100. The three supernatants were combined and Bio-Lyte Ampholyte™ 3/10 (40%) of BioRad was added to a final concentration of 1%. The solubilized proteins were subjected to isoelectric focusing. 60 ml of sample was applied to the Rotofor™ system of BioRad, using 0.1 M phosphoric acid as anode solution, and 0.1 M NaOH as cathode solution. The isoelectric focusing was performed for about 4 hours at 10°C using 10 Watt constant power until the current remained constant for 30 mins. Proteins were divided into two separate groups in accordance with their pI: proteins having a pI in the range of 2-6.5 are referred to as Pool 1 proteins while proteins having a pI in the range of 6.5-10 are referred to as Pool 2 proteins. Pool 1 and Pool 2 proteins were harvested separately and diluted 1:50 with PBS.

Example 3 Injection of Pool 1 and Pool 2 proteins**Method**

The following four preparations were injected intradermally into the forearm of a schizophrenic patient:

- i. Pool 2 proteins (marked as P2)
- ii. Pool 1 proteins (marked as P1)
- iii. Autologous platelets (marked A)
- iv. PBS.

0.1 ml of each above preparation were injected and the prepara-

tions were injected at four different injection sites spaced about 10 cm from each other. The skin reaction at each injection site was monitored 24 h, 48 h and 72 h after injection.

Results

The results are seen in Fig. 1 where (i) is the highest injection site on the arm and (iv) is the lowest.

As seen in the figure, a DTH response measured as explained above, was observed in the two schizophrenic patients at the site of injection of Pool 2 proteins and no DTH reaction was seen at the site of P1 injection. Furthermore, the DTH reaction at the P2 site of injection was substantially enhanced as compared to the DTH reaction seen at the site of injection of the autologous platelets. Thus, it is, in most cases, preferred to use a P2 protein preparation obtained from a pool of blood samples obtained from several heterologous individuals in the diagnostic assay of the invention.

CLAIMS:

1. A protein preparation comprising platelet derived proteins or fractions thereof having an isoelectric point (pI) above about 6.5 and preferably within the range of above 6.5 to about 9.5, said preparation capable of eliciting a DTH reaction in a schizophrenic individual upon injection thereof to the individual.
2. A diagnostic method for assaying schizophrenia in a subject comprising:
 - a) obtaining a preparation comprising, as an active component, platelet derived proteins or fractions thereof having a pI above about 6.5;
 - b) injecting said preparation into a subject; and
 - c) examining the subject for the occurrence of delayed type hypersensitivity reaction at the site of the injection, a positive result being a reaction above that which is observed in non-schizophrenic subjects, indicating that the subject has a high likelihood of being schizophrenic.
3. A method for the preparation of a reagent for use in diagnosis of schizophrenia, comprising:
 - a) obtaining blood samples from a number of individuals, preparing a pool from said samples and collecting platelets therefrom;
 - b) preparing a protein fraction from said platelet preparation comprising proteins or fractions thereof having a pI of above about 6.5.
4. A diagnostic method for assaying schizophrenia in a subject comprising:
 - (a) obtaining a blood sample from a number of schizophrenic and/or non schizophrenic individuals other than the tested subject and collecting platelets therefrom;

(b) preparing a protein fraction from said platelet separation comprising proteins or fractions thereof having a pI of above about 6.5;

(c) injecting said protein preparation into a subject; and

(d) examining the subject for the occurrence of a delayed type hypersensitivity reaction at the site of the injection, a positive result being a reaction above that which is observed in non-schizophrenic subjects, indicating that the subject has a high likelihood of being schizophrenic.

5. A diagnostic method for assaying schizophrenia in a subject comprising;

a) obtaining a blood sample from an individual and collecting platelets therefrom;

b) collecting proteins or fractions thereof from said platelet sample, said proteins or fractions having a pI of above about 6.5.

c) injecting said collected proteins or fractions thereof to the tested individual; and

d) examining the subject for the occurrence of delayed type hypersensitivity reaction at the site of the injection, a positive result being a reaction above that which is observed in non-schizophrenic subjects, indicating that the subject has a high likelihood of being schizophrenic.

6. The method of any one of the previous claims, wherein said proteins or fractions thereof have a pI within the range of above 6.5 to about 9.5.

7. A kit for use in diagnosis of schizophrenia comprising a protein or a fraction thereof prepared from human platelets, said proteins or fractions thereof having a pI of above about 6.5 together with instructions for use.

8. A kit in accordance with Claim 6, wherein the proteins or fractions thereof have a pI within the range of above 6.5 to about 9.5.

9. A kit in accordance with Claims 6 or 7, wherein the proteins or fractions thereof are prepared from heterologous platelets obtained from a number of individuals other than the individual to be tested.
10. A kit in accordance with Claims 6 or 7, wherein the proteins or fractions thereof are prepared from autologous platelets obtained from the individual to be tested.

P₂

P₁

A

C




Fig. 1



INTERNATIONAL SEARCH REPORT

Internatic Application No
PCT/IL 98/00592

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 13152 A (YEDA RESEARCH AND DEVELOPMENT COMPANY LTD) 10 April 1997 see page 12, line 10 - line 23; claims 1-4	1, 3, 7-10
Y	----	1-10
Y	BIOLOGICAL ABSTRACTS, vol. 82, Philadelphia, PA, US; abstract no. 24088, XP002100216 see abstract & G. SH. BURBAEA ET AL.: "The Arthus reaction and delayed hypersensitivity reaction to neurospecific proteins S-100 and 10-40-4 in schizophrenic patients." ZK NEVROPATOL PSIKHIATR IM S S KORSAKOVA, vol. 86, no. 1, 1986, pages 193-195, Moscow RUS -----	1-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

19 April 1999

03/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bohemen, C

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 98/00592

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9713152 A	10-04-1997	AU 6999296 A	28-04-1997
		CA 2233321 A	10-04-1997
		EP 0856158 A	05-08-1998
<hr/>			